

ORIGINAL ARTICLE

# L-type $\text{Ca}^{2+}$ channel blockade with antihypertensive medication disrupts VTA synaptic plasticity and drug-associated contextual memory

M Degoulet<sup>1</sup>, CE Stelly<sup>1</sup>, K-C Ahn<sup>1,2</sup> and H Morikawa<sup>1</sup>

Drug addiction is driven, in part, by powerful and enduring memories of sensory cues associated with drug intake. As such, relapse to drug use during abstinence is frequently triggered by an encounter with drug-associated cues, including the drug itself. L-type  $\text{Ca}^{2+}$  channels (LTCCs) are known to regulate different forms of synaptic plasticity, the major neural substrate for learning and memory, in various brain areas. Long-term potentiation (LTP) of NMDA receptor (NMDAR)-mediated glutamatergic transmission in the ventral tegmental area (VTA) may contribute to the increased motivational valence of drug-associated cues triggering relapse. In this study, using rat brain slices, we found that isradipine, a general LTCC antagonist used as antihypertensive medication, not only blocks the induction of NMDAR LTP but also promotes the reversal of previously induced LTP in the VTA. In behaving rats, isradipine injected into the VTA suppressed the acquisition of cocaine-paired contextual cue memory assessed using a conditioned place preference (CPP) paradigm. Furthermore, administration of isradipine or a  $\text{Ca}_v1.3$  subtype-selective LTCC antagonist (systemic or intra-VTA) before a single extinction or reinstatement session, while having no immediate effect at the time of administration, abolished previously acquired cocaine and alcohol (ethanol) CPP on subsequent days. Notably, CPP thus extinguished cannot be reinstated by drug re-exposure, even after 2 weeks of withdrawal. These results suggest that LTCC blockade during exposure to drug-associated cues may cause unlearning of the increased valence of those cues, presumably via reversal of glutamatergic synaptic plasticity in the VTA.

*Molecular Psychiatry* advance online publication, 23 June 2015; doi:10.1038/mp.2015.84

## INTRODUCTION

Addiction is a chronic, relapsing disorder driven in part by strong associations formed between drugs and sensory cues experienced during drug intake, such as places, people and interoceptive drug cues, that are, subjective effects caused by drugs themselves.<sup>1–3</sup> Addictive drugs are thought to hijack synaptic plasticity mechanisms in key brain circuits involved in reward learning, especially the mesolimbic dopaminergic system, comprising the ventral tegmental area (VTA) and its projections to the nucleus accumbens and other limbic structures.<sup>4–6</sup> As such, powerful and enduring memories of drug-related cues are formed, overshadowing other cues associated with non-drug rewards and driving continued drug use as well as relapse after a period of abstinence. Therefore, reducing the strength of drug cue memories by manipulating the underlying synaptic plasticity mechanisms has received particular attention.

During cue-reward conditioning, dopamine neuron burst responses (2–10 action potentials (APs) at 10–50 Hz) shift in time from the reward to the cue. As a consequence, the reward-associated cue acquires positive valence and triggers approach behavior.<sup>7</sup> Glutamatergic inputs activating NMDA receptors (NMDARs) have a critical role in driving burst firing,<sup>8–11</sup> whereas the role of AMPA receptors (AMPA) in burst generation remains controversial.<sup>12,13</sup> In addition to different forms of synaptic plasticity of AMPARs in dopamine neurons,<sup>5,6</sup> NMDAR-mediated

transmission also undergoes long-term potentiation (LTP) following repeated pairing of glutamatergic input stimulation with postsynaptic burst firing,<sup>14</sup> an activity pattern that may be experienced during cue-reward pairing.<sup>15</sup> Hence, this form of glutamatergic synaptic plasticity may contribute, at least partially, to the acquisition of cue-induced burst responses. Induction of LTP requires AP-evoked  $\text{Ca}^{2+}$  signals amplified by preceding activation of metabotropic glutamate receptors (mGluRs, more specifically mGluR1), in addition to the activation of NMDARs themselves, presumably at the glutamatergic inputs to be potentiated.<sup>16</sup> In contrast, previously induced LTP can be reversed when potentiated inputs are repeatedly stimulated in the absence of postsynaptic APs, raising the possibility that cue memory, or learned valence of the cue, could be unlearned under certain conditions.

Voltage-gated  $\text{Ca}^{2+}$  channels are the major source of activity-dependent  $\text{Ca}^{2+}$  influx. Dihydropyridine-sensitive L-type  $\text{Ca}^{2+}$  channels (LTCCs) are a well-established target for antihypertensive medication because of their involvement in excitation-contraction coupling in the cardiovascular system.<sup>17</sup> LTCCs are also widely expressed in the central nervous system and regulate diverse neuronal processes, such as gene expression, cell survival and synaptic plasticity.<sup>18</sup> Dopamine neurons in both the VTA and substantia nigra express LTCCs.<sup>19,20</sup> In the substantia nigra, these channels, particularly the low-threshold  $\text{Ca}_v1.3$  subtype, have been implicated in driving tonic pacemaker firing and, more

<sup>1</sup>Department of Neuroscience and Waggoner Center for Alcohol and Addiction Research, University of Texas at Austin, Austin, TX, USA and <sup>2</sup>Brain Research Centre, Department of Psychiatry, University of British Columbia, Vancouver, BC, Canada. Correspondence: Professor H Morikawa, Department of Neuroscience and Waggoner Center for Alcohol and Addiction Research, University of Texas at Austin, 2400 Speedway, Austin, TX 78712, USA.

E-mail: morikawa@utexas.edu

Received 2 November 2014; revised 14 April 2015; accepted 30 April 2015

recently, in neuronal death associated with Parkinson's disease,<sup>20–22</sup> however, the pathophysiological roles of LTCCs in the VTA remain unclear. A number of studies have reported that systemic administration of LTCC antagonists blocks the acquisition of drug-induced conditioned place preference (CPP),<sup>23–26</sup> a form of Pavlovian contextual cue learning dependent on NMDAR-mediated transmission in the VTA<sup>9,27–29</sup> (but also see ref. 30). Our previous study has shown that acquisition of psychostimulant CPP is inhibited by mGluR1 or NMDAR antagonist in the VTA, whereas CPP expression is attenuated by NMDAR antagonist, but not by mGluR1 antagonist, in the VTA,<sup>31</sup> supporting the potential contribution of NMDAR LTP in driving CPP. Here, mGluR1/NMDAR blockade would suppress CPP acquisition by inhibiting LTP induction at glutamatergic inputs activated by contextual cues of the CPP box, whereas blocking potentiated NMDAR-mediated excitation at those inputs would interfere with CPP expression. In this study, we examined how LTCC blockade in the VTA affects NMDAR LTP in *ex vivo* brain slices and drug (cocaine/ethanol)-induced CPP in behaving rats.

## MATERIALS AND METHODS

### Animals

Male Sprague-Dawley rats (3–10 weeks old; Harlan Laboratories, Houston, TX, USA) were housed in groups of three and maintained on a 12 h light/dark cycle with food and water available *ad libitum*. All animal procedures were approved by the University of Texas Institutional Animal Care and Use Committee.

### Electrophysiology

Horizontal midbrain slices (~200  $\mu\text{m}$ ) were prepared from rats (3–7 weeks old) and recordings were made at 33–35 °C in physiological saline, as in our previous studies.<sup>14,16,31</sup> Recordings were performed in the lateral VTA located 50–150  $\mu\text{m}$  from the medial border of the medial terminal nucleus of the accessory optic tract. Internal solution contained (in mM): 115 K-methylsulfate, 20 KCl, 1.5  $\text{MgCl}_2$ , 10 HEPES, 0.025 EGTA, 2 Mg-ATP, 0.2  $\text{Na}_2\text{-GTP}$  and 10  $\text{Na}_2\text{-phosphocreatine}$  (pH ~7.25, ~285 mOsm  $\text{kg}^{-1}$ ). Putative dopamine neurons were identified by spontaneous firing (1–5 Hz) with broad APs (>1.2 ms) in cell-attached configuration and large  $I_h$  currents (>200 pA; evoked by a 1.5 s hyperpolarizing step of 50 mV) in whole-cell configuration. Voltage-clamp recordings were made at a holding potential of -62 mV, corrected for a liquid junction potential of -7 mV. Recordings were discarded if the series resistance increased above 16 M $\Omega$  or the input resistance dropped below 200 M $\Omega$ .

A 2 ms depolarizing pulse of 55 mV was used to elicit an unclamped AP. Time integral of the outward tail current, termed  $I_{K(\text{Ca})}$ , was calculated between 20 ms and 400–600 ms after the depolarizing pulse (expressed in pC).  $I_{K(\text{Ca})}$  thus measured is eliminated by tetrodotoxin and by apamin, a selective antagonist of  $\text{Ca}^{2+}$ -activated small-conductance calcium-activated potassium channels, and thus can be used as a readout of AP-induced  $\text{Ca}^{2+}$  transients.<sup>16</sup>

Loose-patch recordings (~20 M $\Omega$  seal) were made using pipettes filled with 150 mM NaCl to monitor dopamine neuron firing. Aspartate iontophoresis (1 M L-aspartate in ~100 M $\Omega$  pipette placed at ~10–50  $\mu\text{m}$  from the soma/proximal dendrites) was used to evoke NMDAR-dependent bursts.<sup>10,16</sup> Amplitude (~100–200 nA) and duration (~50–150 ms) of the iontophoretic current was adjusted to produce a burst of 5–10 spikes with a minimum instantaneous frequency of 15 Hz.

### NMDAR LTP experiments

Synaptic stimuli were applied every 30 s using a bipolar tungsten electrode (~300  $\mu\text{m}$  tip separation) placed rostral to the recorded neuron. To isolate NMDAR excitatory postsynaptic currents (EPSCs), recordings were performed in the presence of 6,7-dinitroquinoxaline-2,3-dione (10  $\mu\text{M}$ ), picrotoxin (100  $\mu\text{M}$ ), CGP55845 (50 nM), and eticlopride (100 nM) to block AMPA/kainate, GABA<sub>A</sub>, GABA<sub>B</sub> and D<sub>2</sub> dopamine receptors, and in glycine (20  $\mu\text{M}$ ) and low  $\text{Mg}^{2+}$  (0.1 mM) to enhance NMDAR activation. Stimulation intensity was adjusted after break-in (within ~1 min) to obtain ~100 pA EPSCs. Cells with baseline EPSC amplitude (averaged from 10 traces during 5 min window before LTP induction) outside the 90–110 pA range were excluded.

Following 10 min baseline EPSC recording, the effect of sustained synaptic stimulation (33 stimuli at 33 Hz) on  $I_{K(\text{Ca})}$  was assessed immediately before LTP induction. Here,  $I_{K(\text{Ca})}$  was evoked by a single AP alone and an AP with preceding synaptic stimulation (140 ms interval between the offset of synaptic stimulation and AP; each repeated twice). LTP was induced by pairing sustained synaptic stimulation (50 stimuli at 33 Hz) with a burst (five APs at 20 Hz), where the burst onset was delayed by 1 s from the onset of synaptic stimulation. This synaptic stimulation-burst pairing was repeated 10 times every 20 s. In LTP reversal experiments, sustained synaptic stimulation alone or synaptic stimulation paired with a single AP (delayed by 1 s from the synaptic stimulation onset) was delivered repeatedly (10 or 30 times) 30 min after LTP induction. Magnitude of LTP and its reversal was determined by averaged EPSC amplitude from a 5 min window (10 traces) immediately before LTP induction and that from 5 min windows at 25–30 min after LTP induction/reversal. For AP5 experiments (Figure 3c), a 5 min window before AP5 perfusion (that is, 20–25 min after LTP induction) was used.

### Place conditioning

A CPP box (Med Associates) consisting of two distinct compartments separated by a small middle chamber was used for conditioning. Rats (4–10 weeks old) were first subjected to a pretest, in which they explored the entire CPP box for 15 min. The percentage of time spent in each compartment was determined after excluding the time spent in the middle chamber. Rats with initial side preference >60% were excluded. During the next 6 days, rats were given saline injection (1 ml  $\text{kg}^{-1}$ ) and confined to one compartment (days 1, 3, 5) or received cocaine injection (10-mg  $\text{kg}^{-1}$ , intraperitoneal (i.p.)) and confined to the other compartment (days 2, 4 and 6; 15 min each). For ethanol CPP, rats were given saline (4.2 ml  $\text{kg}^{-1}$ ) or ethanol (0.5 g  $\text{kg}^{-1}$ , 15% v/v, i.p.) injection and confined to one compartment for 7 min. Compartment assignment was counterbalanced such that animals had, on average, ~50% initial preference for the drug-paired side in the pretest. A 15 min posttest was performed 1 day after the last conditioning session. In extinction experiments, animals underwent repeated posttests (once daily). For reinstatement, rats received priming injection of cocaine (10 mg  $\text{kg}^{-1}$ ) or ethanol (0.5 g  $\text{kg}^{-1}$ ) prior to the posttest. In some CPP experiments, rats received bilateral intra-VTA infusion (0.3  $\mu\text{l}$  per side, 0.15  $\mu\text{l min}^{-1}$ ) of (1) isradipine (0.6 pmol) or vehicle (0.01% ethanol (=1.7 mM)), (2) compound 8 (6 pmol) or vehicle (0.02% dimethyl sulfoxide) or (3) AP5 (0.6 nmol) or vehicle (phosphate-buffered saline). Intra-VTA microinjection procedure is detailed in Supplementary Materials and Methods. Data from rats with injection sites outside the VTA were excluded from the analysis.


### Data analysis

No statistical methods were used to predetermine sample sizes, but our sample sizes are similar to those reported in previous publications.<sup>14,16,31–33</sup> Group assignment was mostly done in a random fashion, except for certain CPP experiments (Figure 4 and 5; Supplementary Figures S11 and S12), where rats were assigned to treatment groups in a counterbalanced manner based on the first posttest data. Data acquisition and analysis were not blinded. Data are expressed as mean  $\pm$  s.e.m. with the sample size in each group indicated. Data distribution was assumed to be normal, but this was not formally tested. Statistical significance was determined by two-tailed *t*-test or ANOVA followed by Bonferroni *post hoc* test using GraphPad Prism (significant at  $P < 0.05$ ; details provided in figure legends).

## RESULTS

### Isradipine inhibits induction of NMDAR LTP

To gain insight into the LTCC-dependent mechanisms in the VTA, we performed electrophysiological recordings in *ex vivo* VTA slices to examine the effects of isradipine, a dihydropyridine LTCC antagonist used as antihypertensive medication in humans. Isradipine was first tested on NMDAR-dependent dopamine neuron excitation/bursting, which likely has an important role in the acquisition of CPP, as well as its expression.<sup>9,27–29,31</sup> Bath application of isradipine (2  $\mu\text{M}$ ) had no effect on NMDAR-mediated EPSCs elicited by local synaptic stimulation (Figure 1a) or on NMDAR-dependent burst firing evoked by aspartate iontophoresis<sup>10</sup> (Figure 1b). Furthermore, isradipine had no effect




**Figure 1.** Isradipine blocks NMDAR LTP induction in the VTA. **(a)** Isradipine ( $2 \mu\text{M}$ ) had no effect on NMDAR excitatory postsynaptic currents (EPSCs) ( $n=6$  cells; example EPSC traces before and after isradipine application). **(b)** Example traces (top; aspartate iontophoresis at arrows) and summary time graphs (bottom) illustrating that isradipine had no effect on the frequency/number of spikes within the burst ( $n=5$  cells) or tonic firing ( $n=8$  cells). **(c)** Example experiments (EPSC traces at the times indicated) and summary time graph showing that isradipine blocked the induction of NMDAR LTP. Graph at the bottom right depicts average EPSC amplitude during baseline and after LTP ( $F_{1,25} = 21.89$ ,  $P < 0.001$ ,  $n = 12-15$  cells per group; mixed two-way ANOVA).  $***P < 0.001$  vs baseline;  $###P < 0.001$  between groups (Bonferroni *post hoc* test). **(d)** Isradipine blocked LTP induction without affecting synaptic facilitation of  $I_{K(Ca)}$ . Data are from the same cells shown in **(c)** (LTP:  $t_{25} = 4.71$ ,  $P < 0.001$ ;  $I_{K(Ca)}$  facilitation:  $t_{25} = 0.79$ ,  $P = 0.43$ ; unpaired *t*-test). Example traces depict  $I_{K(Ca)}$  evoked by a single AP alone and with preceding synaptic stimulation. Bottom graph illustrates the relationship between LTP magnitude and the degree of  $I_{K(Ca)}$  facilitation obtained from each cell (dashed lines: linear fit to all data points in each group).

on tonic firing (Figure 1b), consistent with an LTCC-independent mechanism of pacemaker activity of dopamine neurons in the VTA.<sup>20,21,34</sup>

Next, we asked if isradipine interferes with the mGluR1-dependent induction of NMDAR LTP. LTP was induced using a synaptic stimulation-burst pairing protocol (see Materials and

Methods). Here, sustained glutamatergic input stimulation leads to mGluR1-dependent production of the cytosolic messenger inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ), which amplifies AP-evoked  $\text{Ca}^{2+}$  signals via  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release from intracellular stores.<sup>14</sup> Baseline NMDAR EPSC amplitude, recorded in low  $\text{Mg}^{2+}$  ( $0.1 \text{ mM}$ ), was set at  $\sim 100 \text{ pA}$  to control for synaptic stimulation intensity and thus for



**Figure 2.** Isradipine in the VTA blocks cocaine CPP acquisition. **(a and b)** Timeline of experiments (top; systemic (i.p.) or intra-VTA injection of isradipine (ISR)/vehicle (VEH) made at arrows) and summary graphs depicting changes in the preference for the cocaine (COC)-paired compartment after three conditioning sessions **(a)** systemic:  $F_{1,30} = 41.08$ ,  $P < 0.001$ ,  $n = 16$  rats per group; intra-VTA:  $F_{1,12} = 26.71$ ,  $P < 0.001$ ,  $n = 6-8$  rats per group; **(b)**  $F_{1,12} = 17.92$ ,  $P < 0.01$ ,  $n = 7$  rats per group; mixed two-way ANOVA). \* $P < 0.05$ , \*\*\* $P < 0.001$  vs pretest; ### $P < 0.001$  between groups (Bonferroni *post hoc* test).

the degree of synaptic activation of the mGluR-IP<sub>3</sub> pathway. Under these conditions, repeated synaptic stimulation-burst pairing (10 times) produced LTP of NMDAR EPSCs that gradually developed over ~30 min (Figure 1c). As in previous studies,<sup>14,32,33</sup> LTP magnitude was positively correlated with the degree of synaptic mGluR-induced facilitation of AP-evoked Ca<sup>2+</sup> signals (assessed immediately before LTP induction using the size of SK currents, termed  $I_{K(Ca)}$ ) (Figure 1d). In contrast, LTP was virtually abolished when isradipine was applied 5 min before and during the delivery of LTP induction protocol, although synaptic facilitation of  $I_{K(Ca)}$  at the time of induction was comparable to that observed in control solution. In separate experiments, we confirmed that isradipine, which did suppress Ca<sup>2+</sup> currents evoked by small depolarizations (10–15 mV) from –62 mV (Supplementary Figure S1), had no effect on the size of burst-evoked  $I_{K(Ca)}$  or the magnitude of  $I_{K(Ca)}$  facilitation produced by photolytic application of IP<sub>3</sub> (Supplementary Figure S2). Therefore, LTCC inhibition with isradipine suppresses LTP induction without affecting burst-evoked Ca<sup>2+</sup> signals or the mGluR/IP<sub>3</sub>-dependent amplification machinery (for example, the size of IP<sub>3</sub>-sensitive Ca<sup>2+</sup> stores (Supplementary Figure S3)).

We further examined if enhancing LTCC activation with the LTCC agonist S(-)-Bay K 8644 promotes NMDAR LTP induction. The magnitude of LTP induced in the presence of Bay K 8644 (1 μM) was comparable to that in control (Supplementary Figure S4). Furthermore, inhibiting NMDARs with AP5 blocked LTP induced in Bay K 8644, as has been reported for LTP in control solution.<sup>14</sup> Hence, increasing LTCC activation appears to have no significant effect on NMDAR LTP induction, in contrast to the major suppression of LTP observed with LTCC inhibition.

#### LTCC blockade in the VTA inhibits acquisition of cocaine CPP

Systemic injection (i.p.) of isradipine has been shown to suppress the acquisition of psychostimulant (cocaine and amphetamine) CPP.<sup>23,24</sup> Blockade of NMDAR LTP induction in the VTA might contribute to CPP suppression. Thus, we sought to determine whether isradipine affects acquisition of cocaine CPP via its effect in the VTA. We found that bilateral intra-VTA injection of isradipine (0.6 pmol per 0.3 μl (= 2 μM) in each side) 5 min before each of the three cocaine conditioning sessions completely blocked CPP acquisition, as was observed with systemic isradipine injection (1.2 mg kg<sup>-1</sup>, i.p.) made 10 min prior to each conditioning session (Figure 2a). Isradipine (both systemic and intra-VTA) had no


significant effect on the overall activity during the conditioning sessions (Supplementary Figure S5). Furthermore, CPP was partially suppressed when intra-VTA isradipine injection was made immediately after each cocaine conditioning session (Figure 2b) (note that the burst-inducing effect of cocaine (10 mg kg<sup>-1</sup>, i.p.) persists >45 min<sup>35</sup> and thus should be still robust after the 15 min conditioning session). Intra-VTA injection of isradipine by itself did not affect side preference (Supplementary Figure S6). These results are consistent with the involvement of LTCC-dependent plasticity processes in the VTA in the acquisition of appetitive cocaine cue memory and, likely, its consolidation.

Next we examined if cocaine conditioning alters NMDAR-mediated excitation in the VTA. In these experiments, we used rats that had undergone cocaine conditioning with systemic injection of isradipine or vehicle (Figure 2a, left panel; recordings made 1 day after the posttest), and the data were compared with those from control rats with no cocaine conditioning experience. There was no change in overall NMDAR-dependent excitation assessed with bath application of NMDA (10 μM; Supplementary Figure S7a), as has been reported.<sup>36,37</sup> A recent study in mice has shown that *in vivo* cocaine experience induces synaptic insertion of GluN3A-containing NMDARs, which display reduced Mg<sup>2+</sup> blockade at hyperpolarized potentials.<sup>38</sup> However, NMDAR EPSCs measured in normal Mg<sup>2+</sup> (1.2 mM) displayed similar voltage dependence in the three groups of rats (Supplementary Figure S7b). Finally, comparable NMDAR LTP magnitude was observed in these groups (Supplementary Figure S7c), as opposed to marked alterations in synaptic plasticity of AMPARs produced by cocaine experience.<sup>36–41</sup> Altogether, these data suggest that cocaine conditioning, with or without isradipine, caused no significant changes in global NMDAR-mediated excitation in the VTA (see Discussion for further details on this issue).

#### Isradipine promotes reversal of NMDAR LTP

Previously induced NMDAR LTP can be reversed by repeated stimulation of the potentiated inputs without postsynaptic firing,<sup>14</sup> which may resemble the activity pattern during extinction training where the cue is repeatedly presented without the reward/drug.<sup>7</sup> Indeed, 30 min after inducing LTP, repeated delivery of the synaptic stimulation train (30 times) (SS-30×; Figure 3a and Supplementary Figure S8a) caused persistent depression of potentiated EPSC amplitude toward the baseline level. In contrast, only transient depression was observed and






**Figure 3.** Isradipine and compound 8 promote reversal of previously induced NMDAR LTP. **(a)** After LTP induction and its development, LTP reversal protocol consisting of (1) sustained synaptic stimulation alone or (2) synaptic stimulation paired with a single AP was repeatedly delivered (30 times; arrowhead). In the third and fourth groups, the latter protocol ('SS+AP-30x') was delivered in the presence of isradipine (ISR) or compound 8 (C8). **(b)** The 'SS+AP-30x' protocol was delivered in control solution or in isradipine without prior LTP induction. **(c)** LTP reversal protocol consisting of synaptic stimulation alone was repeatedly delivered (10 times) in control solution, in isradipine, or in isradipine and AP5 (5  $\mu$ M; produced  $83 \pm 4\%$  peak inhibition of potentiated excitatory postsynaptic currents (EPSCs),  $n = 6$  cells). Summary time graphs of these experiments are shown on the left, whereas graphs depicting average EPSC amplitude during baseline, after LTP (except for **(b)**), and following delivery of LTP reversal protocol are shown on the right (**(a)**  $F_{6,40} = 4.85$ ,  $P < 0.001$ ,  $n = 5-7$  cells/group; **(b)**  $F_{1,10} = 0.01$ ,  $P = 0.94$ ,  $n = 6$  cells per group; **(c)**  $F_{4,32} = 10.49$ ,  $P < 0.001$ ,  $n = 6-7$  cells per group; mixed two-way ANOVA). Example traces for the experiments indicated are shown in the middle. Synaptic stimulation intensity was adjusted to evoke  $\sim 100$  pA baseline EPSCs in each cell; thus the degree of synaptic facilitation of  $I_{K(Ca)}$  was similar in different groups (Supplementary Figures S8d and f).  $**P < 0.01$ ,  $***P < 0.001$  between two LTP stages;  $###P < 0.001$  between groups (Bonferroni *post hoc* test).

EPSC amplitude returned to the LTP level in  $\sim 5-10$  min when (1) a single AP was paired with the synaptic stimulation train (SS+AP-30x; Figure 3a and Supplementary Figure S8a) or (2) the number of repetition of the synaptic stimulation train was reduced to 10 times (SS-10x; Figure 3c and Supplementary Figure S8c). We found that delivery of these protocols in the presence of isradipine invariably caused persistent depression of potentiated EPSCs.

Notably, delivery of the 'SS+AP-30x' protocol in isradipine failed to significantly depress baseline EPSCs without prior LTP induction (Figure 3b and Supplementary Figure S8b). These data demonstrate that isradipine facilitates the reversal, or depotentiation, of previously induced LTP. Application of AP5, which inhibited potentiated NMDAR EPSCs, together with isradipine during the delivery of the 'SS-10x' protocol prevented LTP depotentiation



**Figure 4.** Isradipine and compound 8 promote extinction of cocaine/ethanol CPP and prevent future reinstatement. **(a and b)** Summary graphs depicting the effects of systemic isradipine administration (i.p.) on the expression and extinction of CPP previously induced with cocaine **(a)** or ethanol **(b)** (three conditioning sessions for both). A single injection of isradipine ( $1.2 \text{ mg kg}^{-1}$ ) or vehicle ( $1 \text{ ml kg}^{-1}$  of 16% ethanol ( $0.13 \text{ g kg}^{-1}$ )) was made prior to the second posttest, whereas cocaine/ethanol injections were made immediately before the fifth and seventh posttests to trigger reinstatement (2-week interval between fifth and sixth posttests) **(a)**  $F_{7,126} = 3.40$ ,  $P < 0.001$ ,  $n = 10$  rats per group; **(b)**  $F_{7,91} = 3.21$ ,  $P < 0.01$ ,  $n = 7-8$  rats per group; mixed two-way ANOVA). **(c and d)** Summary graphs showing the effects of intra-VTA injection of isradipine **(c)** or compound 8 **(d)** made before the second posttest following cocaine CPP acquisition. Cocaine-induced reinstatement was tested on the fifth posttest **(c)**  $F_{5,80} = 13.82$ ,  $P < 0.001$ ,  $n = 9$  rats per group; **(d)**  $F_{5,70} = 9.62$ ,  $P < 0.001$ ,  $n = 7-9$  rats per group; mixed two-way ANOVA). **(e)** Intra-VTA isradipine injection had no effect when done immediately after the second posttest ( $F_{3,42} = 0.45$ ,  $P = 0.72$ ,  $n = 7-9$  rats per group; mixed two-way ANOVA). **(f)** Systemic isradipine injection (i.p.) followed by intra-VTA injection of AP5 or vehicle was made before the second posttest ( $F_{3,30} = 9.71$ ,  $P < 0.001$ ,  $n = 6$  rats per group; mixed two-way ANOVA). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  vs pretest; + $P < 0.05$ , +++ $P < 0.001$  between two successive posttests; # $P < 0.05$ , ## $P < 0.01$ , ### $P < 0.001$  between groups (Bonferroni *post hoc* test).

(Figure 3c and Supplementary Figure S8c). Therefore, NMDAR activation at the potentiated inputs is likely required to reverse LTP.

$\text{Ca}_v1.3$  is the major subtype of LTCCs expressed in dopamine neurons.<sup>19,0</sup> A  $\text{Ca}_v1.3$ -selective LTCC antagonist 1-(3-chlorophenethyl)-3-cyclopentylpyrimidine-2,4,6-(1*H*,3*H*,5*H*)-trione, termed compound 8, was recently developed.<sup>42</sup> Indeed, compound 8 ( $20 \mu\text{M}$ ) also enabled the 'SS+AP-30x' protocol to cause depotentiation of potentiated EPSCs (Figure 3a and Supplementary Figure S8a).


#### LTCC blockade in the VTA promotes extinction of cocaine and ethanol CPP

Previously acquired drug CPP gradually diminishes, that is, undergoes extinction, with repeated expression tests (posttests; once daily), where rats are exposed to CPP compartments without the drug. To test if LTCCs are involved in the expression and extinction of cocaine CPP, a single systemic injection of isradipine ( $1.2 \text{ mg kg}^{-1}$ , i.p.) was made 10 min before the second posttest. Isradipine failed to affect CPP expression on the day of injection, in line with the lack of effect of isradipine on tonic and burst firing in VTA slices from rats that had undergone cocaine conditioning (Supplementary Figure S9). However, no significant CPP was observed on the following 2 days (third and fourth posttests, performed without isradipine injection), whereas the vehicle-injected control rats still displayed robust CPP (Figure 4a).

Importantly, priming injection of cocaine before subsequent posttests, which significantly increased CPP in control rats, failed to reinstate CPP previously extinguished in the presence of isradipine, even after 2 weeks of withdrawal in the home cage (that is, no exposure to the CPP box or cocaine). Systemic isradipine injection also promoted extinction of CPP acquired with alcohol (ethanol), a different class of addictive drug, in a similar manner and prevented subsequent ethanol-induced reinstatement (Figure 4b).

Next, we wished to determine whether LTCC blockade in the VTA affects CPP extinction. In cocaine-conditioned animals, we made bilateral intra-VTA injection of isradipine or compound 8 ( $6 \text{ pmol}$  per  $0.3 \mu\text{l}$  ( $= 20 \mu\text{M}$ ) in each side) 5 min before the second posttest, which produced no immediate effect on CPP expression. However, as with systemic isradipine injection, CPP was virtually abolished on subsequent days, and CPP thus extinguished was resistant to cocaine-induced reinstatement (Figures 4c and d). Isradipine/compound 8 injection (systemic or intra-VTA) had no effect on the overall activity during the posttest (Supplementary Figure S10).

Certain manipulations disrupt previously acquired CPP when performed shortly after the posttest, via interacting with the memory reconsolidation process,<sup>43-45</sup> or even without a posttest, via non-specific memory ablation.<sup>46,47</sup> However, intra-VTA injection of isradipine immediately after the second posttest or without a posttest, that is, while rats stayed in the home cage, had no effect on CPP expression the following day (Figure 4e and



**Figure 5.** Isradipine prevents future reinstatement when administered before the posttest following complete extinction. **(a)** Average time course of cocaine CPP extinction during repeated posttests over 9 days ( $F_{9,288} = 77.30$ ,  $P < 0.001$ ,  $n = 33$  rats; repeated measures one-way ANOVA). **(b and c)** A single systemic injection of isradipine or vehicle was made before the tenth posttest performed with **(b)** or without **(c)** cocaine injection (orange arrow), whereas the eleventh posttest was always done with cocaine injection (**(b)**  $F_{2,32} = 16.54$ ,  $P < 0.001$ ,  $n = 9$  rats per group; **(c)**  $F_{2,26} = 30.33$ ,  $P < 0.001$ ,  $n = 7-8$  rats per group; mixed two-way ANOVA).  $**P < 0.01$ ,  $***P < 0.001$  vs pretest;  $+++P < 0.001$  between two successive posttests;  $###P < 0.001$  between groups (Bonferroni *post hoc* test).

Supplementary Figure S11). Thus, to promote CPP extinction, isradipine needs to be present in the VTA during the posttest when rats are exposed to cocaine-associated contextual cues. Accordingly, intra-VTA injection of the NMDAR antagonist AP5, which suppressed CPP expression likely by blocking NMDARs at glutamatergic inputs activated by cocaine-associated cues, prevented CPP extinction induced by systemic isradipine injection (Figure 4f and Supplementary Figure S12).

Repeated posttests over 8–9 consecutive days, during which rats are repeatedly exposed to the CPP compartments without cocaine, resulted in complete extinction of cocaine CPP (Figure 5a). In contrast to extinction induced in the presence of isradipine, CPP simply extinguished with repeated posttests was robustly reinstated by priming injection of cocaine (Figure 5b). However, systemic injection of isradipine, made 10 min before the cocaine-induced reinstatement session (Figure 5b) or before the tenth posttest without cocaine injection (Figure 5c), led to suppression of cocaine-induced reinstatement on the following day.

Altogether, these results show that LTCC blockade during exposure to cocaine/ethanol-paired contextual cues, and to interoceptive cocaine cues during cocaine-induced reinstatement, may cause persistent disruption of appetitive cue memory.

## DISCUSSION

Isradipine, a dihydropyridine LTCC antagonist that crosses the blood–brain barrier, is currently undergoing clinical trials to test if daily isradipine slows neurodegeneration in Parkinson’s disease.<sup>48,49</sup> Our study suggests that isradipine may also be used to treat a critical component of addiction, that is, increased motivational valence of drug-associated cues triggering craving and relapse.

LTCCs are involved in the induction of synaptic plasticity in different brain areas, such as the hippocampus, cerebral cortex and striatum,<sup>50–53</sup> where LTCC activation associated with post-synaptic depolarization is thought to drive synaptic plasticity. In VTA dopamine neurons, our data suggest that basal  $Ca^{2+}$  levels maintained by constant LTCC-mediated  $Ca^{2+}$  influx are essential not only for the induction of NMDAR LTP but also for its maintenance, that is, in preventing LTP reversal triggered by glutamatergic input activity. Here, LTCCs do not contribute to burst-evoked  $Ca^{2+}$  signals or the mGluR/IP<sub>3</sub>-dependent amplification mechanism necessary for LTP induction.<sup>14</sup>  $Ca_v1.3$  LTCCs, which activate at relatively hyperpolarized membrane potentials,<sup>18</sup> are suited for providing tonic  $Ca^{2+}$  influx, even though they do not drive subthreshold oscillations underlying

pacemaker activity in VTA dopamine neurons.<sup>20,21,34</sup> The cellular machinery sensing basal  $Ca^{2+}$  levels (LTCC-dependent) together with transient AP/burst-evoked  $Ca^{2+}$  signals (LTCC-independent) during the delivery of LTP induction and depotentiation protocols remains to be determined. Protein kinase C, which mediates the induction of mGluR/ $Ca^{2+}$ -dependent LTP of NMDAR EPSCs at hippocampal mossy fiber synapses,<sup>54,55</sup> has been ruled out in dopamine neurons.<sup>14</sup>

In the lateral amygdala, LTCC inhibition by antagonists suppresses the induction of AMPAR LTP and impairs aversive Pavlovian conditioning (that is, fear conditioning),<sup>56,57</sup> in a manner analogous to the effects of isradipine in the VTA on NMDAR LTP induction and CPP acquisition. Interestingly, LTCC inhibition in the lateral amygdala during extinction training blocks the extinction of conditioned responses,<sup>58</sup> presumably by interfering with the induction of certain forms of synaptic plasticity within the lateral amygdala underlying inhibitory learning.<sup>59</sup> In contrast, LTCC inhibition in the VTA facilitates the reversal of NMDAR LTP. This may cause unlearning of cue memory by reversing the synaptic plasticity induced during CPP acquisition, thereby promoting CPP extinction and preventing future reinstatement. Hence, LTCC blockade timed with cue exposure would allow for the manipulation of specific cue memory by controlling LTP induction and reversal.<sup>60</sup>

In the present study, no global alterations in NMDAR-dependent excitation were found in the VTA after cocaine conditioning. It remains to be determined whether NMDAR potentiation can be observed specifically at those inputs activated by cocaine-associated cues, as has been demonstrated recently in the lateral amygdala following fear conditioning, where only those inputs paired with a foot shock during conditioning display AMPAR potentiation.<sup>60</sup> Therefore, firm evidence for the role of NMDAR LTP in CPP, or more generally in reward-associated cue learning, is lacking at the moment. Interestingly, a reduction in unitary NMDAR EPSCs at individual glutamatergic synapses has been reported after *in vivo* cocaine experience.<sup>39</sup> This may represent redistribution of NMDARs from synaptic to extrasynaptic sites even if the overall number of NMDARs (synaptic and extrasynaptic) is not changed, as assessed with bath application of NMDA.<sup>61</sup> Alternatively, although speculative, this might represent a form of homeostatic synaptic plasticity,<sup>62</sup> in which NMDAR transmission at inputs not activated by cocaine-associated cues (for example, interoceptive cocaine cues or the experimenter performing injection<sup>40</sup>) are scaled down in response to LTP induced in the presumably small subset of glutamatergic inputs encoding cocaine cues, thereby maintaining the overall strength of NMDAR-dependent excitatory transmission in each neuron. It should be

noted these two possibilities are not necessarily mutually exclusive.

It has been shown that NMDAR blockade in the VTA inhibits the induction of NMDAR LTP and acquisition of psychostimulant CPP.<sup>14,31</sup> In the present study, we found that NMDAR blockade in the VTA also prevents the reversal of NMDAR LTP and extinction of cocaine CPP enabled by isradipine. These results are consistent with the idea that NMDAR activation at specific glutamatergic inputs activated by contextual cues of the CPP box is required for both the learning and unlearning of those cues. In this regard, it is of note that isradipine was effective in preventing future CPP reinstatement, that is, in disrupting cue memory, even when administered before the tenth posttest without CPP expression, where CPP had been completely extinguished during repeated posttests. This suggests that contextual cue inputs were still active during the tenth posttest, thus enabling isradipine to reverse NMDAR LTP at those inputs, but were not capable of supporting CPP expression as a consequence of inhibitory learning during extinction training that would suppress the learned response.<sup>43,44,63</sup> Isradipine administration prior to cocaine-induced reinstatement might further reverse LTP at glutamatergic inputs activated by interoceptive cocaine cues.

Mouse studies with genetic deletion of NMDARs selectively in dopamine neurons reported impaired drug (cocaine/nicotine) and food CPP,<sup>9,27,28</sup> whereas another study observed normal cocaine CPP with impaired reinstatement of extinguished CPP.<sup>30</sup> Although this discrepancy may be due to differences in the CPP protocol (for example, the number and duration of conditioning sessions), these studies overall support the role of NMDARs in dopamine neurons in CPP acquisition, expression and/or reinstatement.

Isradipine, administered systemically or into the VTA, failed to affect the expression of cocaine CPP, including cocaine priming-induced reinstatement of extinguished CPP. This is in line with the lack of effect of isradipine on NMDAR-dependent excitation in the VTA that drives CPP expression.<sup>29,31</sup> Dopamine D1 receptor-mediated activation of LTCCs in the nucleus accumbens has been implicated in the reinstatement of cocaine self-administration.<sup>64</sup> Thus, dopamine regulation of LTCCs in the nucleus accumbens appears to be selectively involved in the expression of operant, but not Pavlovian, drug-seeking behavior.

Current cue exposure-based strategies to treat addiction are aimed at facilitating inhibitory extinction learning (for example, with the NMDAR partial agonist  $\alpha$ -cycloserine) or disrupting memory reconsolidation following retrieval (for example, with the  $\beta$ -adrenergic receptor antagonist propranolol).<sup>43,44</sup> On the basis of rodent studies demonstrating the effectiveness of isradipine and other LTCC antagonists on the acquisition of CPP induced by cocaine and other addictive drugs,<sup>23–26</sup> high dosage of isradipine has been tested in human cocaine addicts in a laboratory setting, which failed to reduce measures of cocaine-induced subjective euphoria with no effect on cognitive performance.<sup>65–67</sup> The present study suggests that isradipine, if taken prior to the retrieval of cue memory, as occurs upon an encounter with environmental cues (places, people and so on) or with interoceptive drug cues during a relapse, would enable unlearning of the increased valence of those cues, thus preventing craving and relapse in the future.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGMENTS

We thank Dr D James Surmeier for compound 8 and Dr Kamran Khodakhah for caged IP<sub>3</sub>. We also thank Garry Cooper for helpful discussions on the effect and usage of compound 8. This work was supported by NIH grants DA015687 and AA015521.

## REFERENCES

- Wise RA, Kiyatkin EA. Differentiating the rapid actions of cocaine. *Nat Rev Neurosci* 2011; **12**: 479–484.
- Robinson TE, Yager LM, Cogan ES, Saunders BT. On the motivational properties of reward cues: individual differences. *Neuropharmacology* 2014; **76**(Pt B): 450–459.
- Volkow ND, Wang GJ, Fowler JS, Tomasi D, Telang F, Baler R. Addiction: decreased reward sensitivity and increased expectation sensitivity conspire to overwhelm the brain's control circuit. *Bioessays* 2010; **32**: 748–755.
- Hyman SE, Malenka RC, Nestler EJ. Neural mechanisms of addiction: the role of reward-related learning and memory. *Annu Rev Neurosci* 2006; **29**: 565–598.
- Kauer JA, Malenka RC. Synaptic plasticity and addiction. *Nat Rev Neurosci* 2007; **8**: 844–858.
- Luthi A, Luscher C. Pathological circuit function underlying addiction and anxiety disorders. *Nat Neurosci* 2014; **17**: 1635–1643.
- Schultz W. Predictive reward signal of dopamine neurons. *J Neurophysiol* 1998; **80**: 1–27.
- Overton PG, Clark D. Burst firing in midbrain dopaminergic neurons. *Brain Res Brain Res Rev* 1997; **25**: 312–334.
- Zweifel LS, Parker JG, Lobb CJ, Rainwater A, Wall VZ, Fadok JP et al. Disruption of NMDAR-dependent burst firing by dopamine neurons provides selective assessment of phasic dopamine-dependent behavior. *Proc Natl Acad Sci USA* 2009; **106**: 7281–7288.
- Morikawa H, Khodakhah K, Williams JT. Two intracellular pathways mediate metabotropic glutamate receptor-induced Ca<sup>2+</sup> mobilization in dopamine neurons. *J Neurosci* 2003; **23**: 149–157.
- Wang LP, Li F, Wang D, Xie K, Wang D, Shen X et al. NMDA receptors in dopaminergic neurons are crucial for habit learning. *Neuron* 2011; **72**: 1055–1066.
- Deister CA, Teagarden MA, Wilson CJ, Paladini CA. An intrinsic neuronal oscillator underlies dopaminergic neuron bursting. *J Neurosci* 2009; **29**: 15888–15897.
- Blythe SN, Atherton JF, Bevan MD. Synaptic activation of dendritic AMPA and NMDA receptors generates transient high-frequency firing in substantia nigra dopamine neurons *in vitro*. *J Neurophysiol* 2007; **97**: 2837–2850.
- Harnett MT, Bernier BE, Ahn KC, Morikawa H. Burst-timing-dependent plasticity of NMDA receptor-mediated transmission in midbrain dopamine neurons. *Neuron* 2009; **62**: 826–838.
- Brown J, Bullock D, Grossberg S. How the basal ganglia use parallel excitatory and inhibitory learning pathways to selectively respond to unexpected rewarding cues. *J Neurosci* 1999; **19**: 10502–10511.
- Cui G, Bernier BE, Harnett MT, Morikawa H. Differential regulation of action potential- and metabotropic glutamate receptor-induced Ca<sup>2+</sup> signals by inositol 1,4,5-trisphosphate in dopaminergic neurons. *J Neurosci* 2007; **27**: 4776–4785.
- Zuccotti A, Clementi S, Reinbothe T, Torrente A, Vandael DH, Pirone A. Structural and functional differences between L-type calcium channels: crucial issues for future selective targeting. *Trends Pharmacol Sci* 2011; **32**: 366–375.
- Lipscombe D, Helton TD, Xu W. L-type calcium channels: the low down. *J Neurophysiol* 2004; **92**: 2633–2641.
- Rajadhyaksha A, Husson I, Satpute SS, Kuppenbender KD, Ren JQ, Guerriero RM et al. L-type Ca<sup>2+</sup> channels mediate adaptation of extracellular signal-regulated kinase 1/2 phosphorylation in the ventral tegmental area after chronic amphetamine treatment. *J Neurosci* 2004; **24**: 7464–7476.
- Chan CS, Guzman JN, Ilijic E, Mercer JN, Rick C, Tkatch T et al. 'Rejuvenation' protects neurons in mouse models of Parkinson's disease. *Nature* 2007; **447**: 1081–1086.
- Guzman JN, Sanchez-Padilla J, Wokosin D, Kondapalli J, Ilijic E, Schumacker PT et al. Oxidant stress evoked by pacemaking in dopaminergic neurons is attenuated by DJ-1. *Nature* 2010; **468**: 696–700.
- Branch SY, Sharma R, Beckstead MJ. Aging decreases L-type calcium channel currents and pacemaker firing fidelity in substantia nigra dopamine neurons. *J Neurosci* 2014; **34**: 9310–9318.
- Pani L, Kuzmin A, Martellotta MC, Gessa GL, Fratta W. The calcium antagonist PN 200-110 inhibits the reinforcing properties of cocaine. *Brain Res Bull* 1991; **26**: 445–447.
- Pucilowski O, Plaznik A, Overstreet DH. Isradipine suppresses amphetamine-induced conditioned place preference and locomotor stimulation in the rat. *Neuropsychopharmacology* 1995; **12**: 239–244.
- Biala G, Langwinski R. Effects of calcium channel antagonists on the reinforcing properties of morphine, ethanol and cocaine as measured by place conditioning. *J Physiol Pharmacol* 1996; **47**: 497–502.
- Pucilowski O, Garges PL, Rezvani AH, Hutheson S, Janowsky DS. Verapamil suppresses d-amphetamine-induced place preference conditioning. *Eur J Pharmacol* 1993; **240**: 89–92.
- Wang LP, Li F, Shen X, Tsien JZ. Conditional knockout of NMDA receptors in dopamine neurons prevents nicotine-conditioned place preference. *PLoS One* 2010; **5**: e8616.



- 28 Zweifel LS, Argilli E, Bonci A, Palmiter RD. Role of NMDA receptors in dopamine neurons for plasticity and addictive behaviors. *Neuron* 2008; **59**: 486–496.
- 29 Tzschentke TM. Measuring reward with the conditioned place preference (CPP) paradigm: update of the last decade. *Addict Biol* 2007; **12**: 227–462.
- 30 Engblom D, Bilbao A, Sanchis-Segura C, Dahan L, Perreau-Lenz S, Bolland B *et al*. Glutamate receptors on dopamine neurons control the persistence of cocaine seeking. *Neuron* 2008; **59**: 497–508.
- 31 Whitaker LR, Degoulet M, Morikawa H. Social deprivation enhances VTA synaptic plasticity and drug-induced contextual learning. *Neuron* 2013; **77**: 335–345.
- 32 Ahn KC, Bernier BE, Harnett MT, Morikawa H. IP3 receptor sensitization during *in vivo* amphetamine experience enhances NMDA receptor plasticity in dopamine neurons of the ventral tegmental area. *J Neurosci* 2010; **30**: 6689–6699.
- 33 Bernier BE, Whitaker LR, Morikawa H. Previous ethanol experience enhances synaptic plasticity of NMDA receptors in the ventral tegmental area. *J Neurosci* 2011; **31**: 5205–5212.
- 34 Khaliq ZM, Bean BP. Pacemaking in dopaminergic ventral tegmental area neurons: depolarizing drive from background and voltage-dependent sodium conductances. *J Neurosci* 2010; **30**: 7401–7413.
- 35 Koulchitsky S, De Backer B, Quertemont E, Charlier C, Seutin V. Differential effects of cocaine on dopamine neuron firing in awake and anesthetized rats. *Neuropsychopharmacology* 2012; **37**: 1559–1571.
- 36 Ungless MA, Whistler JL, Malenka RC, Bonci A. Single cocaine exposure *in vivo* induces long-term potentiation in dopamine neurons. *Nature* 2001; **411**: 583–587.
- 37 Borgland SL, Malenka RC, Bonci A. Acute and chronic cocaine-induced potentiation of synaptic strength in the ventral tegmental area: electrophysiological and behavioral correlates in individual rats. *J Neurosci* 2004; **24**: 7482–7490.
- 38 Yuan T, Mameli M, O' Connor EC, Dey PN, Verpelli C, Sala C *et al*. Expression of cocaine-evoked synaptic plasticity by GluN3A-containing NMDA receptors. *Neuron* 2013; **80**: 1025–1038.
- 39 Mameli M, Bellone C, Brown MT, Luscher C. Cocaine inverts rules for synaptic plasticity of glutamate transmission in the ventral tegmental area. *Nat Neurosci* 2011; **14**: 414–416.
- 40 Chen BT, Bowers MS, Martin M, Hopf FW, Guillory AM, Carelli RM *et al*. Cocaine but not natural reward self-administration nor passive cocaine infusion produces persistent LTP in the VTA. *Neuron* 2008; **59**: 288–297.
- 41 Bellone C, Luscher C. Cocaine triggered AMPA receptor redistribution is reversed *in vivo* by mGluR-dependent long-term depression. *Nat Neurosci* 2006; **9**: 636–641.
- 42 Kang S, Cooper G, Dunne SF, Dusel B, Luan CH, Surmeier DJ *et al*. CaV1.3-selective L-type calcium channel antagonists as potential new therapeutics for Parkinson's disease. *Nat Commun* 2012; **3**: 1146.
- 43 Torregrossa MM, Taylor JR. Learning to forget: manipulating extinction and reconsolidation processes to treat addiction. *Psychopharmacology (Berl)* 2013; **226**: 659–672.
- 44 Milton AL, Everitt BJ. The persistence of maladaptive memory: addiction, drug memories and anti-relapse treatments. *Neurosci Biobehav Rev* 2012; **36**: 1119–1139.
- 45 Xue YX, Luo YX, Wu P, Shi HS, Xue LF, Chen C *et al*. A memory retrieval-extinction procedure to prevent drug craving and relapse. *Science* 2012; **336**: 241–245.
- 46 Lee AM, Kanter BR, Wang D, Lim JP, Zou ME, Qiu C *et al*. Prkcz null mice show normal learning and memory. *Nature* 2013; **493**: 416–419.
- 47 Li YQ, Xue YX, He YY, Li FQ, Xue LF, Xu CM *et al*. Inhibition of PKMzeta in nucleus accumbens core abolishes long-term drug reward memory. *J Neurosci* 2011; **31**: 5436–5446.
- 48 Group PS. Phase II safety, tolerability, and dose selection study of isradipine as a potential disease-modifying intervention in early Parkinson's disease (STEADY-PD). *Mov Disord* 2013; **28**: 1823–1831.
- 49 Ilijic E, Guzman JN, Surmeier DJ. The L-type channel antagonist isradipine is neuroprotective in a mouse model of Parkinson's disease. *Neurobiol Dis* 2011; **43**: 364–371.
- 50 Moosmang S, Haider N, Klugbauer N, Adelsberger H, Langwieser N, Muller J *et al*. Role of hippocampal Cav1.2 Ca<sup>2+</sup> channels in NMDA receptor-independent synaptic plasticity and spatial memory. *J Neurosci* 2005; **25**: 9883–9892.
- 51 Seoane A, Massey PV, Keen H, Bashir ZI, Brown MW. L-type voltage-dependent calcium channel antagonists impair perirhinal long-term recognition memory and plasticity processes. *J Neurosci* 2009; **29**: 9534–9544.
- 52 Adermark L, Lovinger DM. Combined activation of L-type Ca<sup>2+</sup> channels and synaptic transmission is sufficient to induce striatal long-term depression. *J Neurosci* 2007; **27**: 6781–6787.
- 53 Wang Z, Kai L, Day M, Ronesi J, Yin HH, Ding J *et al*. Dopaminergic control of corticostriatal long-term synaptic depression in medium spiny neurons is mediated by cholinergic interneurons. *Neuron* 2006; **50**: 443–452.
- 54 Hunt DL, Castillo PE. Synaptic plasticity of NMDA receptors: mechanisms and functional implications. *Curr Opin Neurobiol* 2012; **22**: 496–508.
- 55 Rebola N, Srikanth BN, Mulle C. Activity-dependent synaptic plasticity of NMDA receptors. *J Physiol* 2010; **588**: 93–99.
- 56 Bauer EP, Schafe GE, LeDoux JE. NMDA receptors and L-type voltage-gated calcium channels contribute to long-term potentiation and different components of fear memory formation in the lateral amygdala. *J Neurosci* 2002; **22**: 5239–5249.
- 57 Langwieser N, Christel CJ, Kleppisch T, Hofmann F, Wotjak CT, Moosmang S. Homeostatic switch in hebbian plasticity and fear learning after sustained loss of Cav1.2 calcium channels. *J Neurosci* 2010; **30**: 8367–8375.
- 58 Davis SE, Bauer EP. L-type voltage-gated calcium channels in the basolateral amygdala are necessary for fear extinction. *J Neurosci* 2012; **32**: 13582–13586.
- 59 Pare D, Duvarci S. Amygdala microcircuits mediating fear expression and extinction. *Curr Opin Neurobiol* 2012; **22**: 717–723.
- 60 Nabavi S, Fox R, Proulx CD, Lin JY, Tsiens RY, Malinow R. Engineering a memory with LTD and LTP. *Nature* 2014; **511**: 348–352.
- 61 Hardingham GE, Bading H. Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nat Rev Neurosci* 2010; **11**: 682–696.
- 62 Turrigiano G. Homeostatic synaptic plasticity: local and global mechanisms for stabilizing neuronal function. *Cold Spring Harb Perspect Biol* 2012; **4**: a005736.
- 63 Pan WX, Schmidt R, Wickens JR, Hyland BI. Tripartite mechanism of extinction suggested by dopamine neuron activity and temporal difference model. *J Neurosci* 2008; **28**: 9619–9631.
- 64 Anderson SM, Famous KR, Sadri-Vakili G, Kumaresan V, Schmidt HD, Bass CE *et al*. CaMKII: a biochemical bridge linking accumbens dopamine and glutamate systems in cocaine seeking. *Nat Neurosci* 2008; **11**: 344–353.
- 65 Johnson BA, Javors MA, Lam YW, Wells LT, Tiouririne M, Roache JD *et al*. Kinetic and cardiovascular comparison of immediate-release isradipine and sustained-release isradipine among non-treatment-seeking, cocaine-dependent individuals. *Prog Neuropsychopharmacol Biol Psychiatry* 2005; **29**: 15–20.
- 66 Roache JD, Johnson BA, Ait-Daoud N, Mauldin JB, Thornton JE, Wells LT *et al*. Effects of repeated-dose isradipine on the abuse liability of cocaine. *Exp Clin Psychopharmacol* 2005; **13**: 319–326.
- 67 Johnson BA, Roache JD, Ait-Daoud N, Wallace CL, Wells LT, Wang Y *et al*. Effects of isradipine on cocaine-induced changes in cognitive performance in recently abstinent cocaine-dependent individuals. *Int J Neuropsychopharmacol* 2005; **8**: 549–556.

Supplementary Information accompanies the paper on the Molecular Psychiatry website (<http://www.nature.com/mp>)